



IQI #19 Hip Fracture Mortality Rate

Provider-Level Indicator

Mortality Indicator for Inpatient Medical Conditions

Numerator

Number of deaths (DISP=20) among cases meeting the inclusion and exclusion rules for the denominator.

Denominator

All discharges, age 65 years and older, with principal diagnosis code for hip fracture.

ICD-9-CM hip fracture diagnosis codes:

82000	FX FEMUR INTRCAPS NOS-CL	82019	FX FEMUR INTRCAP NEC-OPN
82001	FX UP FEMUR EPIPHY-CLOS	82020	TROCHANTERIC FX NOS-CLOS
82002	FX FEMUR, MIDCERVIC-CLOS	82021	INTERTROCHANTERIC FX-CL
82003	FX BASE FEMORAL NCK-CLOS	82022	SUBTROCHANTERIC FX-CLOSE
82009	FX FEMUR INTRCAPS NEC-CL	82030	TROCHANTERIC FX NOS-OPEN
82010	FX FEMUR INTRCAP NOS-OPN	82031	INTERTROCHANTERIC FX-OPN
82011	FX UP FEMUR EPIPHY-OPEN	82032	SUBTROCHANTERIC FX-OPEN
82012	FX FEMUR, MIDCERVIC-OPEN	8208	FX NECK OF FEMUR NOS-CL
82013	FX BASE FEMORAL NCK-OPEN	8209	FX NECK OF FEMUR NOS-OPN

Exclude cases:

- with any diagnosis of periprosthetic fracture
- missing discharge disposition (DISP=missing)
- transferring to another short-term hospital (DISP=2)
- MDC 14 (pregnancy, childbirth, and puerperium)

ICD-9-CM periprosthetic fracture diagnosis code:

99644 MECHANICAL COMPLICATION OF INTERNAL ORTHOPEDIC DEVICE, IMPLANT, AND GRAFT, PERI-PROSTHETIC FRACTURE AROUND PROSTHETIC JOINT