

# Transfusion Reaction Rate

## Patient Safety Indicators #26

### Technical Specifications

#### Area-Level Indicator

#### AHRQ Quality Indicators, Version 4.3, August 2011

Version 4.3a is a maintenance release of Version 4.3. The differences between the two versions are:

- Version 4.3a includes Version 29 of the Limited License edition of the 3M™ APR-DRG Grouper. This grouper corrects an issue found in Version 28 of the Limited License edition of the 3M™ APR-DRG Grouper. The grouper is only used with the Inpatient Quality Indicator (IQI) mortality measures. Version 4.3 includes Version 28 of the Limited License edition of the 3M™ APR-DRG Grouper, which was incorrectly assigning a Risk of Mortality (ROM) subclass for cases dated on or after 10/1/10.
- Version 4.3a allows users to calculate area-level indicators for years 2010 and 2011, correcting an issue previously identified in Version 4.3.

All other aspects of the software, including measure specifications, remain the same. Thus this document (related to Version 4.3) remains unchanged.

## Numerator

Discharges 18 years and older or in MDC 14 with ICD-9-CM codes for transfusion reaction in any diagnosis field (principal or secondary) of all medical and surgical discharges defined by specific DRGs or MS-DRGs.

#### ICD-9-CM Transfusion reaction diagnosis codes:

9996	ABO INCOMPATIBILITY REACTION	E8760	MISMATCHED BLOOD IN TRANSFUSION
9997	RH INCOMPATIBILITY REACTION		

See *Patient Safety Indicators Appendices*:

- Appendix B – Medical Discharge DRGs
- Appendix C – Medical Discharge MS-DRGs
- Appendix D – Surgical Discharge DRGs
- Appendix E – Surgical Discharge MS-DRGs

## Denominator

Discharges in the numerator are assigned to the denominator based on the Metro Area<sup>1</sup> or county of the patient residence, not the Metro Area or county of the hospital where the discharge occurred.

---

<sup>1</sup> The term "metropolitan area" (MA) was adopted by the U.S. Census in 1990 and referred collectively to metropolitan statistical areas (MSAs), consolidated metropolitan statistical areas (CMSAs) and primary metropolitan statistical areas (PMSAs). In addition, "area" could refer to either 1) FIPS county, 2) modified FIPS county, 3) 1999

-----  
Exclude cases:

- with missing gender (SEX=missing), age (AGE=missing), quarter (DQTR=missing), year (YEAR=missing) or principal diagnosis (DX1=missing), county (PSTCO=missing)