

Patient Safety Indicator 26 (PSI 26) Transfusion Reaction Rate

June 2017

Area-Level Indicator
Type of Score: Rate

Prepared by:

Agency for Healthcare Research and Quality
U.S. Department of Health and Human Services
www.qualityindicators.ahrq.gov

DESCRIPTION

Transfusion reactions per 100,000 population, ages 18 years and older.

[NOTE: The software provides the rate per population. However, common practice reports the measure as per 100,000 population. The user must multiply the rate obtained from the software by 100,000 to report events per 100,000 population.]

June 2017 1 of 4

AHRQ QI™ ICD-9-CM Specification Version 6.0 PSI 26 Transfusion Reaction Rate www.qualityindicators.ahrq.gov

NUMERATOR

Surgical and medical discharges, for patients ages 18 years and older or MDC 14 (pregnancy, childbirth, and puerperium), with any-listed ICD-9-CM diagnosis codes for transfusion reaction. Surgical and medical discharges are defined by specific MS-DRG codes.

<u>Appendix C - Medical MS-DRGs</u> Appendix E - Surgical MS-DRGs

Transfusion reaction diagnosis codes: (TRANFID)

9996	Abo incompatibility reaction	9997	Rh incompatibility reaction
99960	Abo incompat react nos	99970	Rh incompat reaction nos
99961	Abo incomp/htr nec	99971	Rh incomp/htr nec
99962	Abo incompat/acute htr	99972	Rh incompat/acute htr
99963	Abo incompat/delay htr	99973	Rh incompat/delay htr
99969	Abo incompat reactn nec	99974	Rh incompat reaction nec

June 2017 2 of 4

AHRQ QI™ ICD-9-CM Specification Version 6.0 PSI 26 Transfusion Reaction Rate www.qualityindicators.ahrq.gov

NUMERATOR EXCLUSIONS

Exclude cases:

• with missing gender (SEX=missing), age (AGE=missing), quarter (DQTR=missing), year (YEAR=missing), principal diagnosis (DX1=missing), or county (PSTCO=missing)

June 2017 3 of 4

AHRQ QI™ ICD-9-CM Specification Version 6.0 PSI 26 Transfusion Reaction Rate www.qualityindicators.ahrq.gov

DENOMINATOR

Population ages 18 years and older in metropolitan area† or county. Discharges in the numerator are assigned to the denominator based on the metropolitan area or county of the patient residence, not the metropolitan area or county of the hospital where the discharge occurred.

† The term "metropolitan area" (MA) was adopted by the U.S. Census in 1990 and referred collectively to metropolitan statistical areas (MSAs), consolidated metropolitan statistical areas (CMSAs) and primary metropolitan statistical areas (PMSAs). In addition, "area" could refer to either 1) FIPS county, 2) modified FIPS county, 3) 1999 OMB Metropolitan Statistical Area or 4) 2003 OMB Metropolitan Statistical Area. Micropolitan Statistical Areas are not used in the QI software.

June 2017 4 of 4