

Hip Fracture Mortality Rate Technical Specifications

Inpatient Quality Indicators #19 (IQI #19)

AHRQ Quality Indicators™, Version 4.5, May 2013

Provider-Level Indicator

Type of Score: Rate

Description

In-hospital deaths per 1,000 hospital discharges with hip fracture as a principal diagnosis for patients ages 65 years and older. Excludes periprosthetic fracture discharges, obstetric discharges, and transfers to another hospital.

[NOTE: The software provides the rate per hospital discharge. However, common practice reports the measure as per 1,000 discharges. The user must multiply the rate obtained from the software by 1,000 to report in-hospital deaths per 1,000 hospital discharges.]

Numerator

Number of deaths (DISP=20) among cases meeting the inclusion and exclusion rules for the denominator.

Denominator

Discharges, for patients ages 65 years and older, with a principal ICD-9-CM diagnosis code for hip fracture.

ICD-9-CM Hip fracture diagnosis codes:

82000	FX FEMUR INTRCAPS NOS-CL	82019	FX FEMUR INTRCAP NEC-OPN
82001	FX UP FEMUR EPIPHY-CLOS	82020	TROCHANTERIC FX NOS-CLOS
82002	FX FEMUR, MIDCERVIC-CLOS	82021	INTERTROCHANTERIC FX-CL
82003	FX BASE FEMORAL NCK-CLOS	82022	SUBTROCHANTERIC FX-CLOSE
82009	FX FEMUR INTRCAPS NEC-CL	82030	TROCHANTERIC FX NOS-OPEN
82010	FX FEMUR INTRCAP NOS-OPN	82031	INTERTROCHANTERIC FX-OPN
82011	FX UP FEMUR EPIPHY-OPEN	82032	SUBTROCHANTERIC FX-OPEN
82012	FX FEMUR, MIDCERVIC-OPEN	8208	FX NECK OF FEMUR NOS-CL
82013	FX BASE FEMORAL NCK-OPEN	8209	FX NECK OF FEMUR NOS-OPN

Exclude cases:

- with any-listed ICD-9-CM diagnosis codes for periprosthetic fracture
- transferring to another short-term hospital (DISP=2)
- MDC 14 (pregnancy, childbirth, and puerperium)

- with missing discharge disposition (DISP=missing), gender (SEX=missing), age (AGE=missing), quarter (DQTR=missing), year (YEAR=missing) or principal diagnosis (DX1=missing)

ICD-9-CM Periprosthetic fracture diagnosis code:
99644 PERIPROSTHETIC FX-PROS JT