



PDI #15 Diabetes Short-term Complications Admission Rate

Area-Level Indicator

Numerator

All non-maternal discharges ages 6 to 17 years with ICD-9-CM principal diagnosis code for short-term complications (ketoacidosis, hyperosmolarity, coma).

ICD-9-CM Ketoacidosis, Hyperosmolarity, Coma diagnosis codes:

25010	DM KETO T2, DM CONT	25022	DM W/ HYPROSM T2, DM UNCNT
25011	DM KETO T1, DM CONT	25023	DM W/ HYPROSM T1, DM UNCNT
25012	DM KETO T2, DM UNCONT	25030	DM COMA NEC TYP II, DM CNT
25013	DM KETO T1, DM UNCONT	25031	DM COMA NEC T1, DM CONT
25020	DM W/ HYPROSM T2, DM CONT	25032	DM COMA NEC T2, DM UNCONT
25021	DM W/ HYPROSM T1, DM CONT	25033	DM COMA NEC T1, DM UNCONT

Exclude cases:

- transfer from a hospital (different facility)
- transfer from a Skilled Nursing Facility (SNF) or Intermediate Care Facility (ICF)
- transfer from another health care facility
- MDC 14 (pregnancy, childbirth, and puerperium)

See *Pediatric Quality Indicators Appendices*:

- Appendix J – Admission Codes for Transfers

Denominator

Population ages 6 to 17 in Metro Area¹ or county.

¹ The term "metropolitan area" (MA) was adopted by the U.S. Census in 1990 and referred collectively to metropolitan statistical areas (MSAs), consolidated metropolitan statistical areas (CMSAs) and primary metropolitan statistical areas (PMSAs). In addition, "area" could refer to either 1) FIPS county, 2) modified FIPS county, 3) 1999 OMB Metropolitan Statistical Area or 3) 2003 OMB Metropolitan Statistical Area. Micropolitan Statistical Areas are not used in the QI software.