



Neonatal Quality Indicator 03 (NQI 03) Neonatal Blood Stream Infection Rate

July 2021

Hospital-Level Indicator

Type of Score: Rate

Prepared by:

Agency for Healthcare Research and Quality

U.S. Department of Health and Human Services

www.qualityindicators.ahrq.gov

DESCRIPTION

• Discharges with healthcare-associated bloodstream infection per 1,000 discharges for newborns and outborns with birth weight of 500 grams or more but less than 1,500 grams; with gestational age between 24 and 30 weeks; or with birth weight of 1,500 grams or more and death, an operating room procedure, mechanical ventilation, or transferring from another hospital within two days of birth. Excludes discharges with a length of stay less than 3 days and discharges with a principal diagnosis (or secondary diagnosis present on admission) of sepsis or bacteremia.

[NOTE: The software provides the rate per hospital discharge. However, common practice reports the measure as per 1,000 discharges. The user must multiply the rate obtained from the software by 1,000 to report events per 1,000 hospital discharges.]

NUMERATOR

Discharges, among cases meeting the inclusion and exclusion rules for the denominator, with either:

- any secondary ICD-10-CM diagnosis code for newborn sepsis (**BSI5DX***) or
- any secondary ICD-10-CM diagnosis code for newborn sepsis requiring a separate organism code or bacteremia (**BSI2DX***), and any secondary ICD-10-CM diagnosis code for staphylococcal or gram-negative bacterial infection (**BSI3DX***)

DENOMINATOR

All discharges for newborns and outborns (*Appendix I*) with either:

- a birth weight of 500 to 1,499 grams (Birth Weight Categories 2, 3, 4 and 5) (*Appendix L*)
- any listed ICD-10-CM diagnosis code for gestational age between 24 and 30 weeks (*GESTC2D**, *GESTC3D**, *GESTC4D**, *GESTC5D**)
- a birth weight greater than or equal to 1,500 grams (Birth Weight Category 0, 6, 7, or 8) and any of:
 - o death (DISP=20)
 - o any listed ICD-10-PCS procedure code for operating room procedure (*Appendix A: ORPROC*)
 - o any listed ICD-10-PCS procedure code for mechanical ventilation (*MECHVCD**)
 - o transferring from another health care facility within two days of birth (ATYPE = 4 and POINTOFORIGINUB04 = 6)

[*Appendix A – Operating Room Procedure Codes*](#)

[*Appendix I – Definitions of Neonate, Newborn, Normal Newborn, and Outborn*](#)

[*Appendix L – Low Birth Weight Categories*](#)

DENOMINATOR EXCLUSIONS

Exclude cases:

- with a principal ICD-10-CM diagnosis code (or secondary diagnosis present on admission) for sepsis or bacteremia (*BSI4DX**)
- with a principal ICD-10-CM diagnosis code (or secondary diagnosis present on admission) for staphylococcal or gram-negative bacterial infection (*BSI3DX**)
- with birth weight less than 500 grams (Birth Weight Category 1) (*Appendix L*)
- with length of stay less than 3 days
- MDC 14 (pregnancy, childbirth, and puerperium)
- with an ungroupable DRG (DRG=999)
- with missing gender (SEX=missing), age (AGE=missing), quarter (DQTR=missing), year (YEAR=missing) or principal diagnosis (DX1=missing)

* See below for code list

Gestational age 24 completed weeks diagnosis codes: (GESTC2D)

P0723 Extreme immaturity of newborn,
gestational age 24 completed weeks

Gestational age between 25 to 26 completed weeks diagnosis codes: (GESTC3D)

P0724 Extreme immaturity of newborn,
gestational age 25 completed weeks

P0725 Extreme immaturity of newborn,
gestational age 26 completed weeks

Gestational age between 27 to 28 completed weeks diagnosis codes: (GESTC4D)

P0726 Extreme immaturity of newborn,
gestational age 27 completed weeks

P0731 Preterm newborn, gestational age 28
completed weeks

Gestational age between 29 to 30 completed weeks diagnosis codes: (GESTC5D)

P0732 Preterm newborn, gestational age 29
completed weeks

P0733 Preterm newborn, gestational age 30
completed weeks

Mechanical ventilation procedure codes: (MECHVCD)

5A1935Z Respiratory ventilation, less than 24
consecutive hours

5A1955Z Respiratory ventilation, greater than 96
consecutive hours

5A1945Z Respiratory ventilation, 24-96
consecutive hours

Newborn sepsis diagnosis code: (BSI5DX)

P3619	Sepsis of newborn due to other streptococci	P3639	Sepsis of newborn due to other staphylococci
P362	Sepsis of newborn due to staphylococcus aureus	P364	Sepsis of newborn due to escherichia coli
P3630	Sepsis of newborn due to unspecified staphylococci		

Newborn septicemia or bacteremia diagnosis codes: (BSI2DX)

P368	Other bacterial Sepsis of newborn	R7881	Bacteremia
------	-----------------------------------	-------	------------

Staphylococcal or Gram-negative bacterial infection diagnosis codes: (BSI3DX)

B961	Klebsiella pneumoniae [k. pneumoniae] as the cause of diseases classified elsewhere	B9623	Unspecified shiga toxin-producing Escherichia coli [E. coli] [stec] as the cause of diseases classified elsewhere
B9620	Unspecified escherichia coli [E. coli] as the cause of diseases classified elsewhere	B9629	Other Escherichia coli [E. coli] as the cause of diseases classified elsewhere
B9621	Shiga toxin-producing Escherichia coli [E. coli] [stec] o157 as the cause of diseases classified elsewhere	B965	Pseudomonas (aeruginosa) (mallei) (pseudomallei) as the cause of diseases classified elsewhere
B9622	Other specified shiga toxin-producing Escherichia coli [E. coli] [stec] as the cause of diseases classified elsewhere	B9689	Other specified bacterial agents as the cause of diseases classified elsewhere

Sepsis or bacteremia diagnosis codes: (BSI4DX)

A021	Salmonella Sepsis	A4189	Other specified Sepsis
A227	Anthrax Sepsis	A419	Sepsis, Unspecified organism
A267	Erysipelothrix Sepsis	A427	Actinomycotic Sepsis
A327	Listerial Sepsis	A5486	Gonococcal Sepsis
A400	Sepsis due to streptococcus, group A	B377	Candidal Sepsis
A401	Sepsis due to streptococcus, group B	P360	Sepsis of newborn due to streptococcus, group B
A403	Sepsis due to streptococcus pneumoniae	P3610	Sepsis of newborn due to Unspecified streptococci
A408	Other streptococcal Sepsis	P3619	Sepsis of newborn due to Other streptococci
A409	Streptococcal Sepsis, Unspecified	P362	Sepsis of newborn due to staphylococcus aureus
A4101	Sepsis due to methicillin susceptible staphylococcus aureus	P3630	Sepsis of newborn due to Unspecified staphylococci
A4102	Sepsis due to methicillin resistant staphylococcus aureus	P3639	Sepsis of newborn due to Other staphylococci
A411	Sepsis due to Other specified staphylococcus	P364	Sepsis of newborn due to Escherichia coli
A412	Sepsis due to Unspecified staphylococcus	P365	Sepsis of newborn due to anaerobes
A413	Sepsis due to hemophilus influenzae	P368	Other bacterial Sepsis of newborn
A414	Sepsis due to anaerobes	P369	Bacterial Sepsis of newborn, Unspecified
A4150	Gram-Negative Sepsis, Unspecified	R6520	Severe Sepsis without septic shock
A4151	Sepsis due to Escherichia coli [E. coli]	R6521	Severe Sepsis with septic shock
A4152	Sepsis due to pseudomonas	R7881	Bacteremia
A4153	Sepsis due to serratia	T8112XA	Postprocedural septic shock, initial encounter
A4159	Other gram-negative Sepsis	T8144XA	Sepsis following a procedure, initial encounter
A4181	Sepsis due to enterococcus		