



**AHRQ Quality Indicators™ (AHRQ QI™)
ICD-9-CM and ICD-10-CM/PCS
Specification Enhanced Version 5.0**

**Prevention Quality Indicator 08 (PQI 08)
Heart Failure Admission Rate**

**October 2015
Area-Level Indicator
Type of Score: Rate**

Prepared by:

Agency for Healthcare Research and Quality
U.S. Department of Health and Human Services
540 Gaither Road
Rockville, MD 20850
www.qualityindicators.ahrq.gov

PQI 08 Heart Failure Admission Rate

DESCRIPTION

Admissions with a principal diagnosis of heart failure per 100,000 population, ages 18 years and older. Excludes cardiac procedure admissions, obstetric admissions, and transfers from other institutions.

[NOTE: The software provides the rate per population. However, common practice reports the measure as per 100,000 population. The user must multiply the rate obtained from the software by 100,000 to report admissions per 100,000 population.]

PQI 08 Heart Failure Admission Rate

NUMERATOR
Discharges, for patients ages 18 years and older, with a principal ICD-9-CM or ICD-10-CM diagnosis code for heart failure.
<i>[NOTE: By definition, discharges with a principal diagnosis of heart failure are precluded from an assignment of MDC 14 by grouper software. Thus, obstetric discharges should not be considered in the PQI rate, though the AHRQ QI™ software does not explicitly exclude obstetric cases.]</i>

Heart failure diagnosis codes: (ACSCHFD) (For discharges prior to October 1, 2001)			
ICD-9-CM	Description	ICD-10-CM	Description
39891	RHEUMATIC HEART FAILURE	<i>No codes for ICD-10-CM</i>	
40201	MAL HYPERT HRT DIS W CHF		
40211	BENIGN HYP HRT DIS W CHF		
40291	HYPERTEN HEART DIS W CHF		
40401	MAL HYPER HRT/REN W CHF		
40403	MAL HYP HRT/REN W CHF/RF		
40411	BEN HYPER HRT/REN W CHF		
40413	BEN HYP HRT/REN W CHF/RF		
40491	HYPERT HRT/REN NOS W CHF		
40493	HYP HT/REN NOS W CHF/RF		
4280	CONGESTIVE HEART FAILURE		
4281	LEFT HEART FAILURE		
42820	SYSTOLIC HRT FAILURE NOS		
42821	AC SYSTOLIC HRT FAILURE		
42822	CHR SYSTOLIC HRT FAILURE		
42823	AC ON CHR SYST HRT FAIL		
42830	DIASTOLC HRT FAILURE NOS		
42831	AC DIASTOLIC HRT FAILURE		
42832	CHR DIASTOLIC HRT FAIL		

42833 AC ON CHR DIAST HRT FAIL
 42840 SYST/DIAST HRT FAIL NOS
 42841 AC SYST/DIASTOL HRT FAIL
 42842 CHR SYST/DIASTL HRT FAIL
 42843 AC/CHR SYST/DIA HRT FAIL
 4289 HEART FAILURE NOS

Heart failure diagnosis codes: (ACSCH2D) (For discharges on or after to October 1, 2001)

ICD-9-CM	Description	ICD-10-CM	Description
39891	RHEUMATIC HEART FAILURE	I0981	Rheumatic heart failure
4280	CONGESTIVE HEART FAILURE	I501	Left ventricular failure
4281	LEFT HEART FAILURE	I5020	Unspecified systolic (congestive) heart failure
42820	SYSTOLIC HRT FAILURE NOS	I5021	Acute systolic (congestive) heart failure
42821	AC SYSTOLIC HRT FAILURE	I5022	Chronic systolic (congestive) heart failure
42822	CHR SYSTOLIC HRT FAILURE	I5023	Acute on chronic systolic (congestive) heart failure
42823	AC ON CHR SYST HRT FAIL	I5030	Unspecified diastolic (congestive) heart failure
42830	DIASTOLC HRT FAILURE NOS	I5031	Acute diastolic (congestive) heart failure
42831	AC DIASTOLIC HRT FAILURE	I5032	Chronic diastolic (congestive) heart failure
42832	CHR DIASTOLIC HRT FAIL	I5033	Acute on chronic diastolic (congestive) heart failure
42833	AC ON CHR DIAST HRT FAIL	I5040	Unspecified combined systolic (congestive) and diastolic (congestive) heart failure
42840	SYST/DIAST HRT FAIL NOS	I5041	Acute combined systolic (congestive) and diastolic (congestive) heart failure
42841	AC SYST/DIASTOL HRT FAIL	I5042	Chronic combined systolic (congestive) and diastolic (congestive) heart failure
42842	CHR SYST/DIASTL HRT FAIL	I5043	Acute on chronic combined systolic (congestive) and diastolic (congestive) heart failure
42843	AC/CHR SYST/DIA HRT FAIL	I509	Heart failure, unspecified
4289	HEART FAILURE NOS		

PQI 08 Heart Failure Admission Rate

NUMERATOR EXCLUSIONS

Exclude cases:

- with any-listed ICD-9-CM or ICD-10-PCS procedure codes for cardiac procedure
- transfer from a hospital (different facility)
- transfer from a Skilled Nursing Facility (SNF) or Intermediate Care Facility (ICF)
- transfer from another health care facility
- with missing gender (SEX=missing), age (AGE=missing), quarter (DQTR=missing), year (YEAR=missing), principal diagnosis (DX1=missing), or county (PSTCO=missing)

[Appendix A – Admission Codes for Transfers](#)

[Appendix B – Cardiac Procedure Codes](#)

PQI 08 Heart Failure Admission Rate

DENOMINATOR

Population ages 18 years and older in metropolitan area† or county. Discharges in the numerator are assigned to the denominator based on the metropolitan area or county of the patient residence, not the metropolitan area or county of the hospital where the discharge occurred.

† The term “metropolitan area” (MA) was adopted by the U.S. Census in 1990 and referred collectively to metropolitan statistical areas (MSAs), consolidated metropolitan statistical areas (CMSAs) and primary metropolitan statistical areas (PMSAs). In addition, “area” could refer to either 1) FIPS county, 2) modified FIPS county, 3) 1999 OMB Metropolitan Statistical Area or 4) 2003 OMB Metropolitan Statistical Area. Micropolitan Statistical Areas are not used in the QI software.

PQI 08 Heart Failure Admission Rate

DENOMINATOR EXCLUSIONS

<i>Not Applicable</i>
