



AHRQ Quality Indicators™ (AHRQ QI™) ICD-10-CM/PCS Specification Version 6.0

Prevention Quality Indicator 01 (PQI 01) Diabetes Short-Term Complications Admission Rate

July 2016

Area-Level Indicator

Type of Score: Rate

Prepared by:

Agency for Healthcare Research and Quality

U.S. Department of Health and Human Services

www.qualityindicators.ahrq.gov

DESCRIPTION

Admissions for a principal diagnosis of diabetes with short-term complications (ketoacidosis, hyperosmolarity, or coma) per 100,000 population, ages 18 years and older. Excludes obstetric admissions and transfers from other institutions.

[NOTE: The software provides the rate per population. However, common practice reports the measure as per 100,000 population. The user must multiply the rate obtained from the software by 100,000 to report admissions per 100,000 population.]

NUMERATOR

Discharges, for patients ages 18 years and older, with a principal ICD-10-CM diagnosis code for diabetes short-term complications (ketoacidosis, hyperosmolarity, or coma).

[NOTE: By definition, discharges with a principal diagnosis of diabetes with short-term complications are precluded from an assignment of MDC 14 by grouper software. Thus, obstetric discharges should not be considered in the PQI rate, though the AHRQ QI™ software does not explicitly exclude obstetric cases.]

Diabetes short-term complications diagnosis codes: (ACDIASD)

E1010	Type 1 diabetes mellitus with ketoacidosis without coma	E1100	Type 2 diabetes mellitus with hyperosmolarity without nonketotic hyperglycemic-hyperosmolar coma (NKHHC)
E1011	Type 1 diabetes mellitus with ketoacidosis with coma	E1101	Type 2 diabetes mellitus with hyperosmolarity with coma
E10641	Type 1 diabetes mellitus with hypoglycemia with coma	E11641	Type 2 diabetes mellitus with hypoglycemia with coma
E1065	Type 1 diabetes mellitus with hyperglycemia	E1165	Type 2 diabetes mellitus with hyperglycemia

NUMERATOR EXCLUSIONS

Exclude cases:

- transfer from a hospital (different facility)
- transfer from a Skilled Nursing Facility (SNF) or Intermediate Care Facility (ICF)
- transfer from another health care facility
- with missing gender (SEX=missing), age (AGE=missing), quarter (DQTR=missing), year (YEAR=missing), principal diagnosis (DX1=missing), or county (PSTCO=missing)

[Appendix A – Admission Codes for Transfers](#)

DENOMINATOR

Population ages 18 years and older in the metropolitan area† or county. Discharges in the numerator are assigned to the denominator based on the metropolitan area or county of the patient residence, not the metropolitan area or county of the hospital where the discharge occurred.‡ May be combined with uncontrolled diabetes as a single indicator as a simple sum of the rates to form the Healthy People 2010 indicator (note that the AHRQ QI™ excludes transfers to avoid double-counting cases).

† The term “metropolitan area” (MA) was adopted by the U.S. Census in 1990 and referred collectively to metropolitan statistical areas (MSAs), consolidated metropolitan statistical areas (CMSAs), and primary metropolitan statistical areas (PMSAs). In addition, “area” could refer to either 1) FIPS county, 2) modified FIPS county, 3) 1999 OMB Metropolitan Statistical Area, or 4) 2003 OMB Metropolitan Statistical Area. Micropolitan Statistical Areas are not used in the QI software.

‡ The denominator can be specified with the diabetic population only and calculated with the SAS QI software through the condition-specific denominator at the state-level feature.