



**AHRQ Quality Indicators™ (AHRQ QI™)
ICD-9-CM and ICD-10-CM/PCS
Specification Enhanced Version 5.0**

**Patient Safety Indicators #24 (PSI #24)
Postoperative Wound Dehiscence Rate**

**October 2015
Area-Level Indicator
Type of Score: Rate**

Prepared by:

Agency for Healthcare Research and Quality
U.S. Department of Health and Human Services
540 Gaither Road
Rockville, MD 20850

www.qualityindicators.ahrq.gov

PSI #24 Postoperative Wound Dehiscence Rate

DESCRIPTION

Postoperative reclosures of the abdominal wall per 100,000 population, ages 18 years and older. Excludes cases with an immunocompromised state and obstetric cases.

[NOTE: The software provides the rate per population. However, common practice reports the measure as per 100,000 population. The user must multiply the rate obtained from the software by 100,000 to report events per 100,000 population.]

PSI #24 Postoperative Wound Dehiscence Rate

NUMERATOR			
Discharges, for patients ages 18 years and older, with any-listed ICD-9-CM or ICD-10-PCS procedure codes for reclosure of postoperative disruption of the abdominal wall.			
<i>Reclosure of postoperative disruption of the abdominal wall procedure codes: (RECLOIP)</i>			
ICD-9-CM	Description	ICD-10-PCS	Description
5461	RECLOSURE OF POSTOPERATIVE DISRUPTION OF ABDOMINAL WALL	0WQFXZZ	Repair Abdominal Wall, External Approach

PSI #24 Postoperative Wound Dehiscence Rate

NUMERATOR EXCLUSIONS

Exclude cases:

- with any-listed ICD-9-CM or ICD-10-CM diagnosis codes or any-listed ICD-9-CM or ICD-10-PCS procedure codes for immunocompromised state
- MDC 14 (pregnancy, childbirth, and puerperium)
- with missing gender (SEX=missing), age (AGE=missing), quarter (DQTR=missing), year (YEAR=missing), principal diagnosis (DX1=missing), or county (PSTCO=missing)

[Appendix I – Immunocompromised State Diagnosis and Procedure Codes](#)

PSI #24 Postoperative Wound Dehiscence Rate

DENOMINATOR

Population ages 18 years and older in metropolitan area† or county. Discharges in the numerator are assigned to the denominator based on the metropolitan area or county of the patient residence, not the metropolitan area or county of the hospital where the discharge occurred.

† The term “metropolitan area” (MA) was adopted by the U.S. Census in 1990 and referred collectively to metropolitan statistical areas (MSAs), consolidated metropolitan statistical areas (CMSAs) and primary metropolitan statistical areas (PMSAs). In addition, “area” could refer to either 1) FIPS county, 2) modified FIPS county, 3) 1999 OMB Metropolitan Statistical Area or 4) 2003 OMB Metropolitan Statistical Area. Micropolitan Statistical Areas are not used in the QI software.

PSI #24 Postoperative Wound Dehiscence Rate

DENOMINATOR EXCLUSIONS
<i>Not Applicable</i>